

## GAUSSIAN INTEGER SOLUTIONS FOR THE ELLIPTIC PARABOLOID $x^2 + y^2 = 10z$

Gopalan. M.A<sup>1</sup>, Vidhyalakshmi . S<sup>2</sup>, Shanthi.J<sup>3</sup>

<sup>1,2,3</sup> Department of Mathematics, Shrimati Indira Gandhi college ,  
 Trichy 620002 ,Tamil Nadu , India

mayilgopalan@gmail.com,vidhyasigc@gmail.com, shanthivishvaa@gmail.com

### ABSTRACT

The elliptic paraboloid represented by the ternary quadratic diophantine equation  $x^2 + y^2 = 10z$  is considered and analyzed for its Gaussian integer solutions. Also, knowing a solution, general formulas for generating sequence of Gaussian integer solutions for the above equation are presented.

**Keywords:** Elliptic paraboloid, Ternary quadratic, Gaussian integer.

**Subject classification:** 11D09.

### INTRODUCTION

The ternary quadratic diophantine equations are rich in variety and have been attracted by many Mathematicians [1-24] and the solutions are represented by non-zero distinct integers. In this context, note that an extension of ordinary integers into complex numbers is the Gaussian integers. In [25] Gaussian integer solutions of the pythagorian equation have been presented. The Gaussian integer solutions for the special equations  $z^2 = y^2 + Dx^2$  and  $x^2 + y^2 = 2z^2$  are presented in [26,27] respectively. Also, in [28] the elliptic paraboloid equation  $3x^2 + 2y^2 = 3z$  has been analyzed for Gaussian integer solutions. This has motivated us to search for Gaussian integer solutions of other ternary quadratic diophantine equations. This communication concerns with the problem of obtaining Gaussian integer solutions of the elliptic paraboloid  $x^2 + y^2 = 10z$ . . Also, knowing a solution, general formulas for

generating sequence of Gaussian integer solutions for the above equation are presented.

### METHOD OF ANALYSIS:

Consider the ternary quadratic equation representing the elliptic paraboloid given by

$$x^2 + y^2 = 10z \quad (1)$$

We present below different patterns of Gaussian integer solutions to (1)

#### Pattern 1:

Introducing the linear transformations  $x = 2f + i10, y = -5 + i3f, z = e + if$  (2) in (1), it is written as

$$5f^2 + 10e + 75 = 0$$

which is satisfied by

$$f = 2\alpha + 1, \quad e = -(2\alpha^2 + 2\alpha + 8)$$

Substituting the values of  $e$  &  $f$  in (2), the Gaussian integer solutions to (1) are given by

$$x = (4\alpha + 2) + i10$$

$$y = -5 + i(6\alpha + 3)$$

$$z = -(2\alpha^2 + 2\alpha + 8) + i(2\alpha + 1)$$

#### Pattern 2:

Introducing the transformations

$$x = 3f + i, y = 2f + i, z = e + if \quad (3)$$

in (1), it is written as

$$5f^2 - 10e - 2 = 0$$

which is satisfied by

$$f = \pm 2^{2k-1}, \quad e = \frac{1}{5} [13(2^{4k-3}) - 1]$$

Substituting the values of  $e$  &  $f$  in (3), the Gaussian integer solutions to (1) are given by

$$\begin{aligned} x &= 3(2^{2k-1}) + i \\ y &= 2^{2k} + i \\ z &= \frac{1}{5} [13(2^{4k-3}) - 1] + i[2^{2k} - 1] \end{aligned}$$

**Pattern 3:**

Assuming

$$x = 2f + i25, y = -5 + i9f, z = e + if \quad (4)$$

in (1), it is written as

$$77f^2 + 10e + 600 = 0$$

$$\text{Let } f = 10F$$

$$\text{Then, } e = -770F^2 - 60$$

Substituting the values of  $e$  &  $f$  in (4), the Gaussian integer solutions to (1) are given by

$$x = 20F + i25$$

$$y = -5 + i90F$$

$$z = -770F^2 - 60 + i10F$$

**Pattern 4:**

Introducing the transformation

$$x = 3f + i, y = 1 + i2f, z = (e^2 + 1) + if \quad (5)$$

in (1), it is written as

$$f^2 = 2e^2 + 2 \quad (6)$$

The initial solution of (6) is

$$e_0 = 1, \quad f_0 = 2$$

To find the other solutions of (6), consider the Pellian equation

$$f^2 = 2e^2 + 1 \quad (7)$$

Whose general solution  $(\tilde{e}_n, \tilde{f}_n)$  is given by

$$\tilde{f}_n = \frac{1}{2} [(3 + 2\sqrt{2})^{n+1} + (3 - 2\sqrt{2})^{n+1}]$$

$$\tilde{e}_n = \frac{1}{2\sqrt{2}} [(3 + 2\sqrt{2})^{n+1} - (3 - 2\sqrt{2})^{n+1}]$$

where  $n = -1, 0, 1, \dots$

Applying the Bhramagutha lemma between the solutions  $(e_0, f_0)$  &  $(\tilde{e}_n, \tilde{f}_n)$  the other values of equation (6) are given by

$$\begin{aligned} e_{n+1} &= \tilde{f}_n + 2\tilde{e}_n \\ f_{n+1} &= 2\tilde{f}_n + 2\tilde{e}_n \end{aligned}$$

Substituting the above values in (5), the Gaussian integer solutions to (1) are given by

$$x_{n+1} = 6\tilde{f}_n + 6\tilde{e}_n + i$$

$$y_{n+1} = 1 + i4[\tilde{e}_n + \tilde{f}_n]$$

$$z_{n+1} = \tilde{f}_n^2 + 4\tilde{e}_n^2 + 4\tilde{e}_n\tilde{f}_n + 1 + i2[\tilde{e}_n + \tilde{f}_n]$$

**Pattern 5:**

Consider

$$x = 5A + i5B$$

$$y = 5A + i5D$$

$$z = 2(B^2 + D^2) + i5A(B + D) \quad (8)$$

Using (8) in (1), it is written as

$$10A^2 - 9B^2 = 9D^2 \quad (9)$$

Introducing the linear transformations

$$\begin{aligned} A &= 3X + 9T \\ B &= 3X + 10T \end{aligned} \quad (10)$$

in (9), we have

$$X^2 = 10T^2 + D^2 \quad (11)$$

which is satisfied by

$$T = 2rs$$

$$D = 10r^2 - s^2$$

$$X = 10r^2 + s^2$$

Substituting the values of  $X, T, D$  in (10) and (8), we have

$$x = 5[(30r^2 + 3s^2 + 18rs) + i(30r^2 + 3s^2 + 20rs)]$$

$$y = 5[(30r^2 + 3s^2 + 18rs) + i(10r^2 - s^2)^2]$$

$$z = 2\{(30r^2 + 3s^2 + 20rs)^2 + (10r^2 - s^2)^2\} + i5\{(30r^2 + 3s^2 + 18rs)(40r^2 + 2s^2 + 20rs)\}$$

which represents the Gaussian integer solutions of (1).

**Pattern 6:**

Note that (11) can be represented as the system of double equations in different ways as follows.

**System 1:**

$$X + D = 5T, \quad X - D = 2T$$

**System 2:**

$$X + D = T^2, \quad X - D = 10$$

**System 3:**

$$X + D = 5T^2, \quad X - D = 2$$

Solving each of the above three systems, we obtain the values of  $X, T, D$

Substituting the above values of  $X, T, D$  in (10) and (8), the corresponding Gaussian integer solutions to (1) are obtained and they are represented as follows.

**Solutions for system 1:**

$$\begin{aligned} x &= 5k(39 + i41) \\ y &= 5k(39 + i3) \\ z &= 20k^2(169 + i429) \end{aligned}$$

**Solutions for system 2:**

$$x = 5[(6k^2 + 18k + 15) + i(6k^2 + 20k + 15)]$$

$$y = 5[(6k^2 + 18k + 15) + i(2k^2 - 5)]$$

$$z = 2\{(6k^2 + 20k + 15)^2 + (2k^2 - 5)^2\} + i5\{(6k^2 + 18k + 15)(8k^2 + 20k + 10)\}$$

**Solutions for system 3:**

$$x = 5[(30k^2 + 18k + 3) + i(30k^2 + 20k + 3)]$$

$$y = 5[(30k^2 + 18k + 3) + i(10k^2 - 1)]$$

$$z = 2\{(30k^2 + 20k + 3)^2 + (10k^2 - 1)^2\} + i5\{(30k^2 + 18k + 3)(40k^2 + 20k + 2)\}$$

**Generation of solutions:**

Now, given Gaussian integer solution to (1), can an infinite number of Gaussian integer solutions be generated? Albeit tacitly, the answer for this question is in the affirmative.

**Generation 1:**

Let  $(x_0, y_0, z_0)$  be a Gaussian integer solution of (1).

Let

$$x_1 = h - kx_0, y_1 = h - ky_0, z_1 = k^2z_0 + h \dots\dots (12)$$

be the second Gaussian integer solutions to (1). Here 'k' is any given non-zero integer and 'h' is a non-zero Gaussian integer. Substituting (12) in (1) and simplifying, we have

$$h = k[x_0 + y_0] + 5$$

Using this value of 'h' in (12), we have

$$x_1 = ky_0 + 5, y_1 = kx_0 + 5, z_1 = k[x_0 + y_0] + k^2z_0 + 5$$

Repeating the above process, the general solution  $(x_n, y_n, z_n)$  to (1) is given by

$$x_n = \frac{k^n}{2} \{(1 - (-1)^n)y_0 + (1 + (-1)^n)x_0\} + 5 \frac{(1 - k^n)}{(1 - k)}$$

$$y_n = \frac{k^n}{2} \{(1 - (-1)^n)x_0 + (1 + (-1)^n)y_0\} + 5 \frac{(1 - k^n)}{(1 - k)}$$

$$z_n = k^n \left(\frac{1 - k^n}{1 - k}\right) (x_0 + y_0) + k^{2n}z_0 + 5 \left(\frac{1 - k^n}{1 - k}\right)^2$$

**Generation 2:**

Let  $(x_0, y_0, z_0)$  be a Gaussian integer solution of (1).

The second solution is obtained by setting

$$x_1 = x_0 + 2h, y_1 = y_0 + 2h, z_1 = z_0 + h^2$$

where 'h' is a non-zero Gaussian integer given by

$$h = 2[x_0 + y_0]$$

The repetition of the above process leads to the general form of Gaussian integer solution to (1) given by

$$\begin{aligned} x_n &= [(9^n + 1)x_0 + (9^n - 1)y_0] \\ y_n &= [(9^n - 1)x_0 + (9^n + 1)y_0] \\ z_n &= z_0 + 4(x_0 + y_0)^2 \left(\frac{9^{2n} - 1}{80}\right) \end{aligned}$$

**Conclusion:**

In this paper, we have presented infinitely many Gaussian integer solutions to the elliptic paraboloid given by  $x^2 + y^2 = 10z$ . Also, we have illustrated a method for generating sequence of Gaussian integer solutions for the given equation, knowing its initial solution. As the ternary quadratic equation are rich in variety, one may search for Gaussian integer solutions for other choices of elliptic paraboloids.

## REFERENCES:

1. Dickson LE. History of Theory of Numbers and Diophantine Analysis, Vol2, Dover publications, New York 2005.
2. Mordell LJ. Diophantine Equations” Academic Press, New York 1970.
3. Carmichael RD. The Theory of Numbers and Diophantine Analysis Dover publications., New York 1959.
4. Gopalan MA, Geetha D. Lattice points on the Hyperboloid of two sheets  $x^2 - 6xy + y^2 + 6x - 2y + 5 = z^2 + 4$  Impact J Sci Tech 2010;4:23-32.
5. Gopalan MA, Vidhyalakshmi S, Kavitha A. Integral points on the Homogeneous cone  $z^2 = 2x^2 - 7y^2$ , The Diophantus J Math 2012;1(2):127-136.
6. Gopalan MA, Vidhyalakshmi S, Sumathi .G Lattice points on the Hyperboloid of one sheet  $4z^2 = 2x^2 + 3y^2 - 4$ . Diophantus J Math 2012; 1(2): 109-115.
7. Gopalan MA, Vidhyalakshmi S, Lakshmi K. Integral points on the Hyperboloid of two sheets  $3y^2 = 7x^2 - z^2 + 21$  Diophantus J Math 2012; 1(2): 99-107.
8. Gopalan MA, Vidhyalakshmi S, Mallika.S Observations on Hyperboloid of one sheet  $x^2 + 2y^2 - z^2 = 2$ . Bessel J Math 2012; 2(3): 221-226.
9. Gopalan MA, Vidhyalakshmi S, Usha Rani TR, Mallika S, Integral points on the Homogeneous cone  $6z^2 + 3y^2 - 2x^2 = 0$ , The Impact J Sci Tech 2012;6(1):7-13.
10. Gopalan MA, Vidhyalakshmi S, Sumathi.G Lattice points on the Elliptic paraboloid  $z = 9x^2 + 4y^2$ , Advances in Theoretical and Applied Mathematics 2012;m7(4):379-385
11. Gopalan MA, Vidhyalakshmi S, Usha Rani TR Integral points on the non-homogeneous cone  $2z^2 + 4xy + 8x - 4z = 0$ , Global Journal of Mathematics and Mathematical sciences 2012;2(1):61-67
12. Gopalan MA, Vidhyalakshmi S, Lakshmi K. Lattice points on the Elliptic paraboloid  $16y^2 + 9z^2 = 4x$ , Bessel J of Math 2013; 3(2): 137-145.
13. Gopalan MA, Vidhyalakshmi S, Uma Rani J, Integral points on the Homogeneous cone  $4y^2 + x^2 = 37z^2$ , Cayley J of Math 2013;2(2):101- 107.
14. Gopalan MA, Vidhyalakshmi S, Kavitha A. Observations on the Hyperboloid of two sheets  $7x^2 - 3y^2 = z^2 + z(y - x) + 4$ . International Journal of Latest Research in Science and technology 2013; 2(2): 84-86.
15. Gopalan MA, Sivagami B. Integral points on the homogeneous cone  $z^2 = 3x^2 + 6y^2$ . IOSR Journal of Mathematics 2013; 8(4): 24-29.
16. Gopalan MA, Geetha V. Lattice points on the homogeneous cone  $z^2 = 2x^2 + 8y^2 - 6xy$  Indian journal of Science 2013; 2: 93-96.
17. Gopalan MA, Vidhyalakshmi S, Maheswari D. Integral points on the homogeneous cone  $35z^2 = 2x^2 + 3y^2$ . Indian journal of Science 2014; 7: 6-10.
18. Gopalan MA, Vidhyalakshmi S, Umarani J. On the Ternary Quadratic Diophantine Equation  $6(x^2 + y^2) - 8xy = 21z^2$ . Sch J Eng Tech 2014; 2(2A): 108-112.
19. Meena K, Vidhyalakshmi S, Gopalan MA, Priya IK. Integral points on the cone  $3(x^2 + y^2) - 5xy = 47z^2$  Bulletin of Mathematics and statistic Research 2014; 2(1): 65-70.
20. Gopalan MA, Vidhyalakshmi S, Nivetha S On Ternary Quadratic Diophantine Equation  $4(x^2 + y^2) - 7xy = 31z^2$ . Diophantus J Math 2014; 3(1): 1-7.
21. Meena K, Vidhyalakshmi S, Gopalan MA Thangam SA. Integral solutions on the homogeneous cone  $28z^2 = 4x^2 + 3y^2$ .

- Bulletin of Mathematics and statistic  
Research 2014; 2(1): 47-53.
- 22.Sanathi J ,Gopalan MA, Vidhyalakshmi.  
Lattice points on the homogeneous cone  
 $8(x^2 + y^2) - 15xy = 56z^2$ . Sch Journal of  
Phy Math Stat 2014; 1(1): 29-32.
- 23.Gopalan MA, Vidhyalakshmi S ,Mallika S  
On Ternary Quadratic Diophantine Equation  
 $8(x^2 + y^2) - 15xy = 80z^2$ .BOMSR 2014;  
2(4): 429-433.
- 24.Meena S,Gopalan MA, Vidhyalakshmi S ,  
Thiruniraiselvi N.Observations on the Ternary  
Quadratic Diophantine Equation  
 $x^2 + 9y^2 = 50z^2$ .International Journal of  
Applied Research 2015; 1(2): 51-53.
- 25.M.A.Gopalan, and S. Vidhyalakshmi,  
Pythagorean Triples of Gaussian Integers,  
DiophantusJournal of Mathematics, 2012,  
vol 1(1); 47-50.
- 26.M.A.Gopalan, G. Sangeetha and Manju  
Somanath, Gaussian Integer Solution for a  
special equation  $z^2 = y^2 + Dx^2$ , presented  
in the International Conference held at  
Bishop Heber College , (HICAMS, January  
5-7, 2012)
- 27.M.A.Gopalan , G. Sangeetha and Manju  
Somanath, Gaussian integer Solution for a  
Special Equation  $x^2 + y^2 = 2z^2$ , Advances  
in Theortical and Applied Mathematics,  
2012, vol 7(4), 329-335.
- 28.M.A.Gopalan , G. Sangeetha and Manju  
Somanath, Gaussian integer Solution for a  
Special Elliptic paraboloid equation  
 $3x^2 + 2y^2 = 3z$ , IJMA, Vol-5(2), 2012,  
159-162.