

GC-MS Analysis and Antimicrobial Activity of Sudanese *Pithecellobium dulce* (Roxb.) Benth Fixed Oil

Abdel Karim. M.^{1(*)}, Fath El-Rahman, A.² and W. Ahmed K.³

1. Sudan University of Science and Technology, Faculty of Science, Dept. of Chemistry
2. Omdurman Islamic University, Faculty of Education, Dept. of Chemistry
3. Omdurman Islamic University, Faculty of Pharmacy

Abstract:-

The present study was aimed to quantify and identify the chemical constituents of *Pithecellobium dulce* (Roxb.) Benth seed oil and to evaluate its antimicrobial activity. Twenty one components were detected by GC-MS analysis. Major constituents are: 9,12-octadecadienoic acid(23.59%), 9-octadecenoic acid (22.65%), methyl 20-methylheneicosanoate(13.99%), hexadecanoic acid(12.88%), tetracosanoic acid(7.45%) and methyl stearate(5.78%) . Butylated hydroxytoluene, a potent antioxidant, was detected as a minor constituent(0.17%). The antibacterial activity of the oil was evaluated via cup plate agar diffusion assay against six standard human pathogens(Gram positive: *Staphylococcus aureus* and *Bacillus subtilis*; Gram negative : *Escherichia coli* and *Pseudomonasa aeruginosa* and the fungi *Candida albicans* and *Aspergillus niger*) . The oil showed different antimicrobial responses against test organisms. It gave partial activity against *Bacillus subtilis* and the fungi *Candida albicans* and *Aspergillus niger*. These results indicate that the oil is a candidate for further optimization

Keywords: *Pithecellobium dulce* , Fixed oil, GC-MS, Antimicrobial activity

Introduction

Pithecellobium species are widely distributed through tropics, specially in Asia and America. They belong to the sub-family Mimosoideae in Legume family. *Pithecellobium dulce* (Roxb.) Benth. (Manila Tamarind) is a multipurpose tropical fruit tree used primarily for its fruits, which are eaten fresh or processed and seeds are processed for non-food uses (Lewis *et.al.*2005). *Pithecellobium dulce* is a small-to medium- sized semi-evergreen tree that grows up to 20m height. Crown is spreading but irregular. Trunk is short, about 1 m high, with crooked branches and somewhat shiny branch lets(Grandtner,2005). Manila tamarind originated from a large central American area, stretching from southern California to Colombia and Venezuela. It was successfully planted in small areas in the south Sahelian and north Sudanese zones. It is now widespread (planted and naturalized) in tropical regions (Grandtner,2005). Manila Tamarind is reported to be a folk medicine for an array of human ailments including: leprosy, peptic ulcer, earache and toothache. It is also used by local healers as emollient .Stems are used to cure dysentery, while leaves are used for intestinal disorders. Seeds are claimed to treat ulcers(Chopra *et.al.*,1992; Rajasab and Mahamad,2004; Sivakumar *et.al.*,2005). Seeds have been reported to contain steroids, saponins, lipids, phospholipids, glycosides, glycolipids and polysaccharides. Bark yielded 37% tannins of the catechol type. Quercitin, kaempferol, dulcitol and afezilin were reported from leaves(Lewis *et.al.*2005). Fatty acid analysis of seed extract yielded 9 saturated and 17 unsaturated fatty acids. *Pithecellobium dulce* is a versatile medicinal plant that attracted a worldwide prominence in recent years. All plant parts elaborates a vast array of biologically active phytochemicals and have been demonstrated to exhibit antidiabetic, free radical scavenging, spermicidal, anticonvulsant(Sharma and Mehta,2013), anti-inflammatory(Krishna *et.al.*,1970', arbortacient (Banerjee,2005), antimicrobial(Shanmugakumaran *et.al.*,2005), ,antivenom(Pithayanukul *et.al.*,2005) and protease inhibitory(Delgado *et.al.*,2004) properties .Some interesting secondary metabolites were isolated from this species and some were evaluated for bioactivity(Kulkarni *et.al.*,1992; Nigam *et.al.*,1997; Yoshikawa *et.al.*,1997; Saxena *et.al.*,1998; Niranjana *et.al.*,1998; Saxena VK, Singhal,1999) . The plant is frequently used for bowel movements. The leaves, when applied as plasters is used for pain and venereal sores. Salted decoction of leaves is a treatment for indigestion, it is also used as abortifacient. The bark is used in dysentery, dermatitis and eye inflammation. In Mexico, decoction of leaves is used for earaches, leprosy and toothaches. In

Indian folk medicine, bark is used as astringent, in dysentery and as febrifuge. Also it is used for dermatitis and eye inflammations. Fruits are astringent and barks are claimed to treat ailments ranging from bronchitis, diarrhea, hemorrhages, sores, liver problems to spleen issues (Sugumaran *et.al.*,2008). Due to extensive use of antimicrobial agents in recent years, the antimicrobial resistance became a global issue. Some current antimicrobial agents are now inefficient in controlling diseases (Nascimento *et.al.*,2000; Hay *et.al.*,2005)^{2,3}. To combat this problem measures for exploring novel, save and potent antimicrobial agents should urgently be taken. Plants which were used for centuries in traditional medicine proved to be a rich source for bioactive phytochemicals. Due to their diverse pharmacological properties, plant secondary metabolites (flavonoids, alkaloids...etc) have recently attracted considerable attention. Currently a lot of research is focusing on the bioconstituents of medicinal plants used in traditional medicine. Numerous reports appeared in literature describing the antimicrobial activity of secondary metabolites against a panel of human pathogens (Nascimento *et.al.*,2000; Hay *et.al.*,2005; Sibanda *et.al.*,2007; Indu *et.al.*,2006; Reddy *et.al.*,2007; Sudharameshwari and Radhika, Doughari and Manzara,2008). Hence, the present study was designed to identify and quantify the lipid constituents of *Pithecellobium dulce* seed oil and to evaluate the oil for its antimicrobial activity.

Material and Methods

Materials

Plant material

Seeds of *Pithecellobium dulce* were collected from Khartoum state and authenticated by the Department of Phytochemistry and Taxonomy, National Research Center, Khartoum-Sudan.

Instruments

A Shimadzo GC-MS-QP2010 Ultra instrument with a RTX-5MS column (30m, length ; 0.25mm diameter ; 0.25 µm, thickness) was used.

Test organisms

Pithecellobium dulce oil was screened for antibacterial and antifungal activities using the standard microorganisms shown in table(1).

Table 1: Test organisms

Ser. No	Micro organism	Type
1	<i>Bacillus subtilis</i>	G+ve
2	<i>Staphylococcus aureus</i>	G+ve
3	<i>Pseudomonas aeruginosa</i>	G-ve
4	<i>Escherichia coli</i>	G-ve
5	<i>Aspergillus niger</i>	fungi
6	<i>Candida albicans</i>	fungi

Methods

Extraction of oil from *Pithecellobium dulce* seeds

Powdered shade-dried seeds of *Pithecellobium dulce* (300g) were exhaustively extracted with n-hexane (Soxhlet). The solvent was removed under reduced pressure and the oil was kept in the fridge at 4°C for further manipulation.

Esterification of oil

A Methanolic solution of sodium hydroxide was prepared by dissolving (2g) of sodium hydroxide in (100ml) methanol. A stock solution of methanolic sulphuric acid was prepared by mixing (1ml) of concentrated sulphuric acid with (99ml) methanol.

The oil(2ml) was placed in a test tube and (7ml) of alcoholic sodium hydroxide were added followed by (7ml) of alcoholic sulphuric acid. The tube was stoppered and shaken vigorously for five minutes and then left overnight.(2ml) of supersaturated sodium chloride were added, then (2ml) of normal hexane were added and the tube was vigorously shaken for five minutes .The

Hexane layer was then separated.(5µl) of the hexane extract were mixed with 5ml diethyl ether . The solution was filtered and the filtrate (1µl) was directly injected in the GC-MS vial.

GC-MS analysis

Pithecellobium dulce seed oil was analysed by gas chromatography – mass spectrometry.A Shimadzo GC-MS-QP2010 Ultra instrument with a RTX-5MS column (30m,length ; 0.25mm diameter ; 0.25 µm, thickness)was used. Helium (purity; 99.99 %) was used as carrier gas. Oven temperature program is given in Table 2, while other chromatographic conditions are depicted in Table 3.

Table 2: Oven temperature program

Rate	Temperature(°C)	Hold Time (min. ⁻¹)
-	150.0	1.00
4.00	300.0	0.00

Table 3 : Chromatographic conditions

Column oven temp.	150.0° C
Injection temp.	300.0° C
Injection mode	Split
Flow control mode	Linear velocity
Pressure	139.3KPa
Total flow	50.0ml/min
Column flow	1.54ml/sec.
Linear velocity	47.2cm/sec.
Purge flow	3.0ml/sec.
Split ratio	-1.0

Antimicrobial assay

Preparation of bacterial suspensions

One ml aliquots of 24 hours broth culture of the test organisms were aseptically distributed onto nutrient agar slopes and incubated at 37°C for 24 hours.

The bacterial growth was harvested and washed off with sterile normal saline, and finally suspended in (100 ml) of normal saline to produce a suspension containing about 10^8 - 10^9 colony forming units per ml. The suspension was stored in the refrigerator at 4°C until used. The average number of viable organism per ml of the stock suspension was determined by means of the surface viable counting technique. Serial dilutions of the stock suspension were made in sterile normal saline in tubes and one drop volumes (0.02 ml) of the appropriate dilution was transferred by adjustable volume micropipette onto the surface of dried nutrient agar plates. The plates were allowed to stand for two hours at room temperature for the drop to dry, and then incubated at 37°C for 24 hours.

Preparation of fungal suspensions

Fungal cultures were maintained on dextrose agar incubated at 25°C for four days. The fungal growth was harvested and washed with sterile normal saline, and the suspension was stored in the refrigerator until used.

Testing for antibacterial activity

The cup-plate agar diffusion method was adopted, with some minor modifications, to assess the antibacterial activity of oil. (2ml) of the standardized bacterial stock suspension were mixed with (200ml) of sterile molten nutrient agar which was maintained at 45°C in a water bath. (20 ml) Aliquots of the incubated nutrient agar were distributed into sterile Petri dishes. The agar was left to settle and in each of these plates which were divided into two halves, two cups in each half (10 mm in diameter) were cut using sterile cork borer (No 4). Each one of the halves was designed for a sample. Separate Petri dishes were designed for standard antibacterial chemotherapeutic, (ampicillin and gentamycin).

The agar discs were removed, alternate cups were filled with (0.1 ml) sample using adjustable volume microtiter pipette and allowed to diffuse at room temperature for two hours. The plates were then incubated in the upright position at 37°C for 24 hours.

The diameters of the resultant growth inhibition zones were measured in duplicates and averaged.

Results and discussion

GC-MS analysis of *Pithecellobium dulce* fixed oil

Pithecellobium dulce seeds oil was analyzed by GC-MS and the characterization of the constituents was initially accomplished by comparison with the MS library (NIST) and also confirmed by interpretation of the recorded fragmentation pattern.

The GC-MS analysis revealed the presence of 21 components (Table 4). The typical total ion chromatograms (TIC) of hexane extract are shown in Fig.1.




Fig.1: Total ion chromatograms

Table 4 :The typical total ion chromatograms(TIC)

Peak#	R.Time	Area	Area%	Name
1	11.390	173449	0.17	Butylated Hydroxytoluene 9-

2	15.653	184499	0.18	Hexadecenoic acid, methyl ester, (z) -
3	15.851	12947477	12.88	Hexadecanoic acid, methyl ester
4	16.615	76299	0.08	cis-10-Heptadecenoic acid, methyl ester
5	16.821	136513	0.14	Heptadecanoic acid, methyl ester
6	17.514	23724870	23.59	9,12-Octadecadienoic acid (Z,Z)-, methyl (
7	17.565	22780049	22.65	9-Octadecenoic acid, methyl ester, (E)-
8	17.762	5816472	5.78	Methyl stearate
9	19.154	1676226	1.67	Tridecanedial
10	19.279	1355682	1.35	Oxiraneoctanoic acid, 3-octyl-,methyl ester
11	19.319	3585956	3.57	11-Eicosenoic acid, methyl ester
12	19.516	4354065	4.33	Methyl 18-methylnonadecanoate
13	19.672	390840	0.39	Methyl 15-hydroxy-9,12-octadecadienoate
14	20.339	91979	0.09	Heneicosanoic acid, methyl ester
15	20.432	248071	0.25	Phenol, 2,2'-methylenebis[6-(1,1-dimethyl
16	20.648	96254	0.10	Octadecanoic acid, 2,3-dihydroxypropyl e!
17	20.958	636972	0.63	cis-10-Nonadecenoic acid, methyl ester
18	21.141	14071742	13.99	Methyl 20-methyl-heneicosanoate
19	21.897	594787	0.59	Tricosanoic acid, methyl ester
20	21.947	124058	0.12	Oxiraneoctanoic acid, 3-octyl-, methyl est(
21	22.637	7490660	7.45	Tetracosanoic acid, methyl ester
		100556920	100.00	

Fatty acids constituted the major bulk of the oil and two antioxidants :butylated hydroxytoluene and 2,2'-Methylene-bis-[6-(1,1-dimethylethyl)-4-methyl]phenol were detected as minor constituents;0.17% and 0.25% respectively.Some important constituents are discussed below:

9,12-Octadecadienoic acid methyl ester(23.59%)




Fig. 3: Mass spectrum of 9,12-octadecadienoic acid methyl ester

The EI mass spectrum of 9,12-octadecadienoic acid methyl ester is shown in Fig. 3. The peak at m/z 294, which appeared at R.T. 17.514 in total ion chromatogram, corresponds to $M^+[C_{19}H_{34}O_2]^+$. The peak at m/z 263 corresponds to loss of a methoxyl function.

9-Octadecenoic acid methyl ester(22.65%)




Fig. 4: Mass spectrum of 9-octadecenoic acid methyl ester

The EI mass spectrum of 9-octadecenoic acid methyl ester is shown in Fig.4. The peak at m/z 296, which appeared at R.T. 17.565 in total ion chromatogram, corresponds to $M^+[C_{19}H_{36}O_2]^+$. The peak at m/z 265 corresponds to loss of a methoxyl function.

Hexadecanoic acid methyl ester(12.88%)




Fig. 2: Mass spectrum of hexadecanoic acid methyl ester

The EI mass spectrum of hexadecanoic acid methyl ester is shown in Fig. 2. The peak at m/z 270, which appeared at R.T. 15.851 in total ion chromatogram, corresponds to $M^+[C_{17}H_{34}O_2]^+$. The peak at m/z 239 corresponds to loss of a methoxyl function.

Methyl stearate(5.78%)




Fig. 5: Mass spectrum of methyl stearate

The EI mass spectrum of methyl stearate is shown in Fig. 5. The peak at m/z 298, which appeared at R.T. 17.762 in total ion chromatogram, corresponds to $M^+[C_{19}H_{38}O_2]^+$. The peak at m/z 267 corresponds to loss of a methoxyl function.

Cis-11-Eicosenoic acid methyl ester(3.57%)




Fig. 6: Mass spectrum of Cis-11-Eicosenoic acid methyl ester

The EI mass spectrum of Cis-11-eicosenoic acid methyl ester is shown in Fig. 6. The peak at m/z 324, which appeared at R.T. 19.319 in total ion chromatogram, corresponds to $M^+[C_{21}H_{40}O_2]^+$. The peak at m/z 293 corresponds to loss of a methoxyl function.

Butylated hydroxytoluene(0.17%)




Fig. 7: Mass spectrum of butylated hydroxytoluene

The EI mass spectrum of butylated hydroxytoluene is shown in Fig. 7. The peak at m/z 220, which appeared at R.T. 11.390 in total ion chromatogram, corresponds to $M^+[C_{15}H_{24}O]^+$. The peak at 205 is due loss of a methyl function.

2,2`-Methylene-bis-[6-(1,1-dimethylethyl)-4-methyl]phenol(0.25%)




Fig. 8: Mass spectrum of 2,2`-Methylene-bis-[6-(1,1-dimethylethyl)-4-methyl]phenol

The EI mass spectrum of 2,2`-methylene-bis-[6-(1,1-dimethylethyl)-4-methyl]phenol is shown in Fig. 8. The peak at m/z340, which appeared at R.T.20.432 in total ion chromatogram, corresponds to $M^+[C_{15}H_{24}O]^+$. The peak at m/z325 corresponds to loss of CH_3 .

Antimicrobial activity

The oil was evaluated for antimicrobial potency against standard organisms. The average of the diameters of the growth inhibition zones are shown in Table (5). The results were interpreted in terms of the commonly used terms (<9mm: inative;9-12mm:partially active;13-18mm: active;>18mm:very active). Tables (6) and (7) represent the antimicrobial activity of standard antibacterial and antifungal chemotherapeutic agents against standard bacteria and fungi respectively.

Table 5 : Antibacterial activity of *Pithecellobium dulce* oil :M.D.I.Z (mm)

Drug	Conc.(mg/ml)	Ec	Ps	Sa	Bs	Ca	An
oil	100	-	-	8	11	11	10

Table 6 : Antibacterial activity of standard chemotherapeutic agents :M.D.I.Z (mm)

Drug	Conc. mg/ml	Bs.	Sa.	Ec.	Ps.
Ampicillin	40	15	30	-	-
	20	14	25	-	-

	10	11	15	-	-
Gentamycin	40	25	19	22	21
	20	22	18	18	15
	10	17	14	15	12

Table 7 : Antifungal activity of standard chemotherapeutic agents against standard fungi

Drug	Conc. mg/ml	An.	Ca.
Clotrimazole	30	22	38
	15	17	31
	7.5	16	29

Sa.: *Staphylococcus aureus*
 Ec.: *Escherichia coli*
 Pa.: *Pseudomonas aeruginosa*
 An.: *Aspergillus niger*
 Ca.: *Candida albicans*
 Bs.: *Bacillus subtilis*
 M.D.I.Z: Mean diameter or growth inhibition zone (mm)..

The oil showed partial activity against *Bacillus subtilis* and the fungi *Candida albicans* and *Aspergillus niger*. These results indicate that the oil is a candidate for further optimization.

References

Banerjee, A.(2005), *J. Indian Chem. Soc.*, **82**, 186.

Chopra, R.N., Nayar, S.L., Chopra, I.C.(1992) Glossary of Indian Medicinal Plants, Council of Scientific and Industrial Research, New Delhi , P. 195.

Doughari JH, Manzara S.(2008), *Afr. J. Microbiol Res.*, **2**, 67.

Galla N.(2013), *J. Food Pharm Sci* , **1** ,74.

Delgado, V.F., Lopez, V.H.E., Valdes, R.S., Labra, B., Chagolla, L.A., Lopez, V.E.(2004) *J Agricult Food Chem* , **52** (20), 6115.

Hay, A.D., Thomas, M., Montgomery, A., Wetherell, M., Lovering, A., McNult, C.(2005) *J Antimicrob Chemother* , 56, 146.

Indu, M.N., Hatha, A.M., Abirosh, C., Harsha, U., Vivekanandan, G.(2006), *Braz J Microbiol* , **37**, 153.

Kulkarni ,A.S., Khotpal ,R.R., Saha, S.K., Bhakare, H.A.(1992), *J Pharm Sciences* , **54** (3), 105.

Krishna, P.B., Gupta, M.B., Chittranjan, R.(1970) *Indian J Med Res* , **58**, 724.

Lewis, G., Schrire, B., Mackinder, B. and Lock, M.(2005) "Legumes of the World", Royal Botanic Gardens, Kew.

Misra, G., Nigam, S.K., Singh, S., Nigam, S.G., Saxena, R.C.(1979),*Indian Drugs* , **17** (1), 6.

Nascimento, G.G.F., Locatelli, J., Freitas, P.C., Silva, G.L.(2000), *Braz J Microbiol* , **31**, 247.

Nigam, S.K., Misra, G., Uddin, R., Yoshikawa, K., Kawamoto, M., and Arihara, S.(1997), *Phytochemistry* , **44** (7), 1329.

Niranjan, P.S., Kazuo, K., Zhonghua, J., Sukdeb, B., Basudeb, A., Tamotsu, N.(1999) *J Chem Res* , **32**, 558.

Pithayanukul, P., Ruenraroengsak, P., Bavovada, R., Pakmanee, N., Suttisri, R., Saenoon, S.(2005) ,*J Ethno Pharmacol*, **97** (3): 527.

Rajasab, A.H., Mahamad, I.(2004), *Indian Journal of Traditional Knowledge*, **4**, 419.

Reddy, M.K., Gupta, S.K., Jacob, M.R., Khan, S.I., Daneel, F.(2007), *Planta Med* ,**73** (5), 461.

Sharma,S.,Mehla,B.(2013),*Journal of Medical Pharmaceutical and Allied Sciences*, **5**, 37.

Sugumaran,M.,Vetrichelvan,T.,Darlin-Quine,S.(2008), *Ethnobotanical Leaflets* **12**, 446.

Sibanda, T., Okoh, A.I.(2007) ,*Afr J Biotechnol*, **6** (25), 2886.

Sudharameshwari, K., Radhika, J.(2007), *AJTCAM* , **4** (2), 199.

Sivakumar, A.(2005), *Ancient Science of Life*, **30** (2), 69.

Saxena, V.K., Singhal, M.(1999) ,*Fitoterapia* ,**70** (1):, 98.

Saxena, V.K., Singal, M.(1998), *Fitoterapia* , **69** (4), 305.

Shanmugakumaran, S.D., Amerjothy, S., Balakrishna, K., Kumar, M.S.V. (2005), *Indian Drugs* , **42** (6), 392.

Shanmugakumaraan, S.D., Amerjothy, S., Balakrishna, K.(2008), *Phcog Mag* , **4**, 59.

Yoshikawa, K., Suzaki ,Y., Tanaka ,M., Arihara, S., Nigam, S.K.(1997), *J Nat Prod* , **60** (12), 1269.