

g_u -Semi closed sets in generalized topological spaces

Srija S¹, Jayanthi D²

¹ M Sc Mathematics, Avinashilingam University, Coimbatore, Tamil Nadu, India

² Assistant Professor of mathematics, Avinashilingam University, Coimbatore, Tamil Nadu, India

Abstract: In this paper, we have introduced a new class of sets in generalized topological spaces called g_u -semi closed sets. Also we have investigated some of their basic properties.

Keywords: Generalized topological spaces, g -semi closed sets, g_u -semi-closed sets.

1. Introduction: The concept of generalized topological spaces was introduced and investigated by A. Csaszar[1]. Many g -closed sets like g -pre closed set, g -semi closed set etc., in generalized topological spaces are introduced by him. In this paper, we have introduced a new class of sets in generalized topological spaces called g_u -semi closed sets. Also we have investigated some of their basic properties.

2. Preliminaries

Definition 2.1: [1] Let X be a non empty set and let g be a collection of subsets of X . Then g is called a generalized topology (GT for short) on X if and only if $\emptyset \in g$ and $G_i \in g$ for $i \in I \neq \emptyset$ implies $G = \bigcup_{i \in I} G_i \in g$. The pair (X, g) is called as a generalized topological space (GTS for short) on X . The elements of g are called g -open sets and their complements are called g -closed sets.

We denote the family of all g -closed sets in X by $g(X)$. The generalized closure of a subset S of X , denoted by $c_g(S)$, is the intersection of generalized closed sets including S . And the interior of S , denoted by $i_g(S)$, is the union of generalized open sets contained in S .

Note that $c_g(S) = X - i_g(X-S)$ and $i_g(S) = X - c_g(X-S)$.

Definition 2.2: [2] Let (X, g) be a generalized topological space and $A \subseteq X$. Then A is said to be

- (i) g -semi closed if $i_g(c_g(A)) \subseteq A$
- (ii) g -pre closed if $c_g(i_g(A)) \subseteq A$
- (iii) g - α closed if $c_g(i_g(c_g(A))) \subseteq A$
- (iv) g - β closed if $i_g(c_g(i_g(A))) \subseteq A$
- (v) g -regular closed if $c_g(i_g(A)) = A$

The complement of g -semi closed (resp., g -pre closed, g - α closed, g - β closed, g -regular closed) is said to be g -semi open (resp., g -pre open, g - α open, g - β open, g -regular open).

3. g_u -semi closed sets in generalized topological space

In this section we introduced g_u -semi closed sets in generalized topological spaces and studied some of their basic properties.

Definition 3.1: Let (X, g) be a generalized topological space. Then a non empty subset A is said to be g_u -semi closed if $sc_g(A) \subseteq i_g(c_g(U))$ whenever $A \subseteq U$ and U is g -open.

Example 3.2: Let $X = \{a, b, c\}$ and let $g = \{\emptyset, \{a\}, \{b, c\}, X\}$ then (X, g) is a generalized topological space. Now let $A = \{c\}$ and $U = \{b, c\}$. Then $A \subseteq U$ and $sc_g(A) = A \cup i_g(c_g(A)) = A \cup i_g(c_g(\{c\})) = \{b, c\}$ and $i_g(c_g(U)) = i_g(c_g(\{b, c\})) = \{b, c\}$. Therefore $sc_g(A) \subseteq i_g(c_g(U))$. Therefore A is g_u -semi closed set in (X, g) .

Definition 3.3: Let A be a non empty set in a generalized topological spaces in (X, g) . Then the g -semi interior and g -semi closure of A are defined as

$$si_g(A) = \bigcup \{G/G \text{ is a } g\text{-semi open set and } G \subseteq A\} \text{ and}$$

$$sc_g(A) = \bigcap \{G/G \text{ is a } g\text{-semi closed set and } G \supseteq A\}.$$

It is to be noted that for any set A in (X, g) , we have $(sc_g(A))^c = si_g(A^c)$ and $(si_g(A))^c = sc_g(A^c)$

Theorem 3.4: Every g -closed set in (X, g) is a g_u - semi closed sets in (X, g) but not conversely.

Proof: Let A be g -closed in (X, g) then $c_g(A) = A$. Now let $A \subseteq U$ and U is g -open. Then $sc_g(A) = A \cup i_g(c_g(A)) = A \cup i_g(A) \subseteq U = i_g(U) \subseteq i_g(c_g(U))$. Therefore by hypothesis $sc_g(A) \subseteq i_g(c_g(U))$. This implies A is g_u -semi closed set.

Example 3.5: Let $X = \{a, b, c\}$ and let $g = \{\emptyset, \{a\}, \{b, c\}, X\}$ then (X, g) is a generalized topological space. Now let $A = \{c\}$ and $U = \{b, c\}$. Then $A \subseteq U$ and $sc_g(A) = A \cup i_g(c_g(A)) = A \cup i_g(c_g(\{c\})) = \{b, c\}$ and $i_g(c_g(U)) = i_g(c_g(\{b, c\})) = \{b, c\}$. Therefore $sc_g(A) \subseteq i_g(c_g(U))$. Therefore A is g_u -semi closed set in (X, g) . But $c_g(A) = c_g(\{c\}) = \{b, c\} \neq A$. Hence A is not g -closed in (X, g) .

Remark 3.6: Not every g_u - semi closed set in (X, g) is g -pre closed in (X, g) .

Example 3.7: Let $X = \{a, b, c\}$ and let $g = \{\emptyset, \{a, b\}, \{b, c\}, X\}$. Then (X, g) is a generalized topological space. Now let $A = \{a, b\}$ and $U = \{a, b\}$. Then $A \subseteq U$ and $sc_g(A) = A \cup i_g(c_g(A)) = A \cup i_g(c_g(\{a, b\})) = \{a, b\}$ and $i_g(c_g(U)) = i_g(c_g(\{a, b\})) = \{a, b\}$. Therefore $sc_g(A) \subseteq i_g(c_g(U))$. Therefore A is g_u - semi closed set in (X, g) . But $c_g(i_g(A)) = c_g(i_g(\{a\})) = \{a, b\} \not\subseteq A$. Therefore A is not a g -pre closed set in (X, g) .

Remark 3.8: Not every g_u -semi closed set in (X, g) is g - β closed set in (X, g) .

Example 3.9: Let $X = \{a, b, c, d\}$ and let $g = \{\emptyset, \{c\}, \{a, b\}, \{a, b, c\}, \{a, b, d\}, X\}$. Then (X, g) is a generalized topological space. Now let $A = \{a, b\}$ and $U = \{a, b, d\}$. Then $A \subseteq U$ and $sc_g(A) = A \cup i_g(c_g(A)) = A \cup i_g(c_g(\{a, b\})) = \{a, b, d\}$ and $i_g(c_g(U)) = i_g(c_g(\{a, b, d\})) = \{a, b, d\}$. Therefore $sc_g(A) \subseteq i_g(c_g(U))$. Therefore A is g_u - semi closed set in (X, g) . But $i_g(c_g(i_g(A))) = i_g(c_g(i_g(\{a, b\}))) = \{a, b, d\} \not\subseteq A$. Hence A is not a g - β closed set in (X, g) .

Theorem 3.10: Every g -semi closed set in (X, g) is a g_u - semi closed set in (X, g) but not conversely.

Proof: Let (X, g) be a generalized topological space. Let A be g -semi closed. Then $i_g(c_g(A)) \subseteq A$. Now let $A \subseteq U$ and U be g -open. Then $sc_g(A) = A \cup i_g(c_g(A)) = A \cup A = A \subseteq U = i_g(U) \subseteq i_g(c_g(U))$. Therefore A is g_u -semi closed set.

Example 3.11: Let $X = \{a, b, c\}$ and let $g = \{\emptyset, \{a, b\}, \{b, c\}, X\}$. Then (X, g) is a generalized topological space. Now let $A = \{b\}$ and $U = \{a, b\}$. Then $A \subseteq U$ and $sc_g(A) = A \cup i_g(c_g(A)) = A \cup i_g(c_g(\{b\})) = X \subseteq X$ and $i_g(c_g(U)) = i_g(c_g(\{a, b\})) = X$. Therefore $sc_g(A) \subseteq i_g(c_g(U))$. Therefore A is a g_u - semi closed set in (X, g) . But $i_g(c_g(A)) = i_g(c_g(\{b\})) = X \not\subseteq A$. Therefore A is not a g -semi closed set in (X, g) .

Theorem 3.12: Every g - α closed set in (X, g) is a g_u - semi closed set in (X, g) but not conversely.

Proof: Let (X, g) be a generalized topological space. Let A be g - α closed. Then $c_g(i_g(c_g(A))) \subseteq A$. Now let $A \subseteq U$ and U be g -open. Then $sc_g(A) = A \cup i_g(c_g(A)) \subseteq A \cup c_g(i_g(c_g(A))) \subseteq A \cup A \subseteq A \subseteq U = i_g(U) \subseteq i_g(c_g(U))$. Therefore A is g_u -semi closed set.

Example 3.13: Let $X = \{a, b, c\}$ and let $g = \{\emptyset, \{a, b\}, \{b, c\}, X\}$. Then (X, g) is a generalized topological space. Now let $A = \{b\}$ and $U = \{a, b\}$. Then $A \subseteq U$ and $sc_g(A) = A \cup i_g(c_g(A)) = A \cup i_g(c_g(\{b\})) = X \subseteq X$ and $i_g(c_g(U)) = i_g(c_g(\{a, b\})) = X$. Therefore $sc_g(A) \subseteq i_g(c_g(U))$. Therefore A is g_u - semi closed set (X, g) . But $c_g(i_g(c_g(A))) = c_g(i_g(c_g(\{b\}))) = X \not\subseteq A$. Therefore A is not g - α closed set in (X, g) .

Theorem 3.14: Every g -regular closed set in (X, g) is a g_u - semi closed set in (X, g) but not conversely.

Proof: Let A be g -regular closed set. Then $c_g(i_g(A)) = A$. Now let $A \subseteq U$ and U be g -open. Then $sc_g(A) = A \cup i_g(c_g(A)) = A \cup i_g(c_g(i_g(A))) = A \cup i_g(c_g(i_g(A))) = A \cup i_g(A) \subseteq A \subseteq U = i_g(U) \subseteq i_g(c_g(U))$. Therefore A is g_u -semi closed set.

Example 3.15: Let $X = \{a, b, c\}$ and let $g = \{\emptyset, \{a, b\}, \{b, c\}, X\}$. Then (X, g) is a generalized

topological space. Now let $A = \{b\}$ and $U = \{a, b\}$. Then $A \subseteq U$ and $sc_g(A) = A \cup i_g(c_g(A)) = A \cup i_g(c_g(\{b\})) = X \subseteq X$ and $i_g(c_g(U)) = i_g(c_g(\{a, b\})) = X$. Therefore $sc_g(A) \subseteq i_g(c_g(U))$. Therefore A is g_u -semi closed set in (X, g) . But $c_g(i_g(A)) = c_g(i_g(\{b\})) = \emptyset \neq A$. Therefore A is not g -regular closed set in (X, g) .

Theorem 3.16: If a set A is g_u -semi closed in (X, g) then $sc_g(A)$ - A contains no non empty closed set.

Proof: Suppose that A is a g_u -semi closed set (X, g) . Let F be a closed subset of $sc_g(A)$ - A . Then $F \subseteq A^c$ and hence $A \subseteq F^c$. Since F^c is g -open and A is g_u -semi closed set, $sc_g(A) \subseteq F^c$. This implies $F \subseteq (sc_g(A))^c$. Then by Theorem 3.3 we have $F \subseteq sc_g(A)$. But $sc_g(A) \cap (sc_g(A))^c = \emptyset$. Therefore $F = \emptyset$. Hence the theorem is proved.

Theorem 3.17: In a generalized topological space X , for each $x \in X$, $\{x\}$ is g -closed or its complement $X - \{x\}$ is g_u -semi closed in (X, g) .

Proof: Suppose that $\{x\}$ is not g -closed in (X, g) . Then $X - \{x\}$ is not g -open and the only g -open set containing $X - \{x\}$ is X . Therefore $sc_g(X - \{x\}) \subseteq X$. Therefore $X - \{x\}$ is g_u -semi closed in (X, g) .

Theorem 3.18: If A is g -open and g_u -semi closed then A is semi closed.

Proof: Since A is g -open and g_u -semi closed, we have $A \subseteq X$ and $sc_g(A) \subseteq i_g(c_g(U))$. But $i_g(c_g(A)) \subset sc_g(A)$ as $sc_g(A) = A \cup i_g(c_g(A))$. Then $A = sc_g(A)$. Thus A is semi g -closed.

Theorem 3.19: Then the following two conditions are equivalent in a generalized topological space (X, g) ;

- (i) A is g -open and g_u -semi closed
- (ii) A is g -regular open

Proof: (i) \Rightarrow (ii) By (i) As A is g -open set and g -semi closed, we have $A \subseteq A$ also $i_g(A) = A$. Now $i_g(c_g(A)) \subseteq A \cup i_g(c_g(A)) = sc_g(A) \subseteq A = i_g(A) \subseteq i_g(c_g(A))$. Therefore $i_g(c_g(A)) = A$. Therefore A is g -regular open.

(ii) \Rightarrow (i) Since every g -regular open set is g -open, A is g -open and $i_g(c_g(A)) = A$. As $A \subseteq A$, we have $sc_g(A) = A \cup i_g(c_g(A)) = A \cup A = A = i_g(A) \subseteq i_g(c_g(A))$. Thus A is g_u -semi closed in (X, g) .

References:

- [1] A. Csarzar, Generalized open sets in generalized topologies, Acta Math.Hungar., 96(2002), 351-357.
- [2] A.Csarzar, Generalized open sets in generalized topologies, Acta Math.Hungar., 106(2005),53-66.
- [3] Devi, ; Maki., Balachandran , K. :Semi generalized closed maps and generalized closed maps, Mem. Fac. Kochi univer. A. Math.,14(1993),41-54.
- [4] Maki, H; Devi:R. balachandran, K.: Associated topologies of generalized α closed maps and α - generalized closed maps, Mem. Fac. Kochi univer. A. Math.,15(1994),57-63.