

## **A NOTE ON THE MATHEMATICAL MODEL FOR THE DYNAMICS OF BIRD FLU DISEASE**

**S.A. Egbetade and I.A. Salawu**

Department of Mathematics and Statistics,  
The Polytechnic, Ibadan.

### **Abstract**

In a recent paper [1], a model for the dynamics of bird flu disease was studied and it was shown that the disease free equilibrium of the model is stable. We show in this paper, using a reliable Jacobian matrix and model parameter values taken from [1] that one endemic equilibrium is unstable.

### **INTRODUCTION**

Recently, Aderinto [1] proposed a mathematical model for the dynamics of bird flu disease. The author aimed at using the model to examine human population with respect to the disease and its transmission. In particular, he showed that the disease free equilibrium is stable. In the present paper, we extend his results to endemic equilibrium.

### **2. MATHEMATICAL FORMULATION**

The model is [1]

$$S'(t) = \beta N - \frac{\alpha S(t)A(t)I(t)}{N} - (\delta + d)S(t) + \phi_1 A(t) + \phi_2 I(t) \quad (1)$$

$$A'(t) = \frac{\alpha S(t)A(t)}{N} + (1-r)\mu A(t) - \phi_1 A(t) \quad (2)$$

$$I'(t) = \frac{\alpha S(t)A(t)}{N} - (\delta + d + \varphi_2)I(t) + r\mu A(t) - \varphi_1 A(t) \quad (3)$$

Satisfying  $S(0) = S_0 > 0$ ,  $I(0) = I_0 > 0$ ,  $A(0) = A_0 > 0$ ,  $0 < \alpha < 1$ ,  $0 < \delta < 1$ ,  $0 < r < 1$

where

$S(t)$  = number of susceptible individuals at time  $t$

$A(t)$  = number of asymptomatic infective persons (without symptoms) at time  $t$

$I(t)$  = number of infective persons with onset of symptoms at time  $t$

$N$  = total number of birds in the locality considered

$\alpha$  = transmission rate of infection from birds to human

$\beta$  = average birth rate

$\delta$  = disease induced death rate

$d$  = natural death rate

$r$  = fraction of asymptomatic individual progressing to the class  $I(t)$

$\mu$  = rate of exposed to the disease

$\varphi_1$  = recovered rate of the class  $A(t)$

$\varphi_2$  = recovered rate of the class  $I(t)$

### 3. METHOD OF SOLUTION

For the endemic equilibrium  $(0, 0, 1)$ , the Jacobian matrix is

$$J = \begin{pmatrix} -(\delta + d) & \varphi_1 & \varphi_2 \\ 0 & (1-r)\mu - 1 & 0 \\ \alpha\beta & r\mu - 1 & -(\delta + d) + \varphi_2 \end{pmatrix} \quad (4)$$

and the corresponding characteristics equation is given by

$$\lambda^3 + r_1\lambda^2 - r_2\lambda - r_3 = 0 \tag{5}$$

where

$$r_1 = 2(\delta + d) + \mu(1 - r) + \varphi_1 + \varphi_2 \tag{6}$$

$$r_2 = 2\delta [\mu(1 + r) - d - \varphi_2] - d[2(r + \varphi_1) + d + \varphi_2] - \delta (\delta + \varphi_2^2) + \varphi_2(r\mu + \varphi_1) + \alpha\beta \tag{7}$$

$$r_3 = \delta [\mu(\delta^2 - \varphi_2 - r\delta - r\varphi_2 - 2rd) - \varphi_1 (\delta + 2d - \varphi_2)] - \mu d[d + \varphi_2 + r(d - \varphi_2)] - \varphi_1 d(d - \varphi_2) + \alpha\beta [\mu(1 - r) - \varphi_1] \tag{8}$$

Theorem: If  $r_1 \geq 0$ ,  $r_2 \geq 0$ ,  $r_3 \geq 0$ , then equilibrium (0, 0, 1) is unstable.

Proof: For  $r_1 \geq 0$ ,  $r_2 \geq 0$ ,  $r_3 \geq 0$ , equation (5) has one positive root.

Hence the result.

## CONCLUSION

Using the data in [1],  $r_1 = 1.92$ ,  $r_2 = 0.18$ ,  $r_3 = 0.91$ , it shows that J is unstable.

## REFERENCES

- [1] Aderinto, Y.O. (2008). On a mathematical model for the dynamics of bird u disease, Int. J. Num. Maths., 3(1), 30-41.
- [2] Anderson, R.M. (1991). Infectious disease of humans: dynamics and control. Oxford University Press, New York.
- [3] Anderson, R.M. et.al. (1992). The dynamics of circulating influenza strains conferring partial cross-immunity. Journal of Mathematics Biology, 35, 825 – 842.