

Antimicrobial activities of *Solanum incanum*, *Elettaria cardamomum* and *Zingiber officinale*, used traditionally to treat pathogenic microbes.

Dalal Hussien M. Alkhalifah, Associate Professor Microbiology of Biology Dept., Faculty

of Sciences Princess Nourah bint Abdulrahman University, Riyadh, Kingdom of Saudi Arabia

Introduction

In recent times, there have been increases in antibiotic resistant strains of clinically important pathogens, which have led to the emergence of new bacterial strains that are multi-resistant (WHO, 1978, Aibinu *et al.*, 2003; Aibinu *et al.*, 2004). Multiple drug resistance has developed due to the indiscriminate use of commercial antimicrobial drugs commonly used in the treatment of infectious disease (Davis, 1995). The increasing prevalence of multidrug resistant strains of bacteria and the recent appearance of strains with reduced susceptibility to antibiotics raises the specter of untreatable bacterial infections and adds urgency to the search for new infection-fighting strategies (Sieradzki *et al.* 1999). Such a fact is cause for concern, because of the number of patients in hospitals who have suppressed immunity, and due to new bacterial strains, which are multi-resistant. Consequently, new infections can occur in hospitals resulting in high mortality. The indiscriminate use of antibiotics has led to drug resistance of many bacterial strains. The development of multidrug-resistant pathogens has been related to the occurrence of over and under-dosage of antimicrobials (Vargas *et al.*, 2004; Schelz *et al.*, 2006). The emergency of antibiotic resistance by bacteria has become a medical catastrophe and we may be entering a 'post-antibiotic' era where antibiotics are no longer effective. Development of new microbial compounds for resistant organisms is becoming critically important (Martini and Eloff, 1998). Herbal medicine is the oldest and most tried and tested form of medicine. In a sense it forms the basis of all medicine. It is the original medicine, the mother of all remedies used today. It has been used by all cultures for centuries and is still the main form of medical treatment. Herbal medicine is the most important medicine for the majority of people on the planet, especially those who

cannot afford expensive drugs (McKenna, 1996). *Solanum incanum*, commonly known as bitter garden egg belongs to the family Solanaceae. It is a delicate perennial often cultivated as an annual crop. It is a shrub, growing 1 – 3 m high. The leaves are simple, ovate, elliptic, 2.5 – 12 cm long and 2.5 – 8 cm wide. The fruit is fleshy, less than 3 cm in diameter on wild plants but much larger in cultivated forms. Fruit is spherical, green, often striped with white, turning yellow to orange-brown when ripe (Denston, 1951). *Solanum* species are the most potent plants

against pathogenic microorganisms. *Solanum incanum* (L) is one of the important traditional medicinal plants belongs to Solanaceae family. Antibacterial activity of *Solanum incanum* was studied (Britto and Senthilkumar 2001; Pavitra *et al.*, 2012) and presence of analysis of phytochemicals were also studied (Pavitra *et al.*, 2012). Other *Solanum* species, *S. torvum* (leaf, stem and roots) showed antibacterial and antifungal activity (Bari *et al.*, 2010) and antibacterial activity of *Solanum surattense* whole plant extract (Patil *et al.*, 2009) and leaf extract (Sheeba, 2010) were studied. Analysis, presence of phytochemicals and potent antibacterial activity of leaf, root and seed extracts were studied in *S. nigrum* (Sridhar *et al.*, 2011). Cardamom is a dried fruit of *Elettaria cardamomum* belonging to family Zingiberaceae. It is commonly known as queen of spices for the versatile use in culinary practice. Cardamom is a perennial shrub with thick, fleshy, lateral roots which can grow to a height of eight feet (Kapoor, 2000). Soriful *et al.*, (2010) studied the antimicrobial activity of *Elettaria cardamomum* on some Gram positive bacteria: *Staphylococcus aureus*, *Bacillus subtilis*, *Bacillus megaterium*, and *Sarcina lutea* as well as Gram negative bacteria: *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Shigella dysenteriae*, and *Shigella sonnei*. Methanolic extract inhibited the growth of all the tested bacteria having various degrees of inhibition zones. Ginger known as rhizomes of *Zingiber officinale* (Zingiberaceae) is one of the spices valued for its aroma and pungency characteristic. It grows in the tropical region, especially in the southern and eastern part of Asia. That spice is commercialized in the dry form. Ginger is a stimulant, carminative and frequently used for dyspepsia, gastroparesis,

constipation and colic (Wood, 2013). Ginger oil has been proved to prevent skin cancer in mice (Oyagbemi *et al*, 2010). Some studies have demonstrated that gingerols can be used to fight against ovarian cancer (Park *et al*, 2008; McGee, 2004). [10]-gingerol and [12]-gingerol isolated from ginger rhizome have been reported to show antibacterial activity against periodontal bacteria (Bartley and Jacobs, 2000). Previous phytochemical investigations on the rhizome of *Zingiber officinale* yielded 6-gingerol, zingerone, shogaol, butyl hydroxyl toluene, butyl hydroxyl anisole (Imadia *et al*, 1983; Miri *et al*, 2008). The aim of the present study was detection of antimicrobial activities and toxicity of *Solanum incanum*, *Elettaria cardamomum* and *Zingiber officinale*.

Materials and Methods

Plant materials

Aerial parts, fruits, roots and leaves of some plants were collected from local markets in Jeddah, Saudi Arabia during summer 2012. The collected plant materials were put in clean plastic bags and transferred directly to the lab. The plants under investigation were *Solanum incanum*, *Elettaria cardamomum* and *Zingiber officinale*.

Tested organisms

Bacterial and fungal isolates were obtained from Biology Department, Faculty of Science, King Abdul-Aziz University (KAU), Jeddah, Saudi Arabia. The bacterial strains were *Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853, *Klebsiella A. baumannii* ATCC 1656-2, *P. mirabilis* ATCC ,700603, *pneumoniae* (ESBL) ATCC ATCC) 12453, *Staphylococcus aureus* ATCC 25923, *Staphylococcus aureus* (MRSA , and *Enterococcus faecalis* (VRE) ATCC 51299. The tested fungi were 33591 *Aspergillus flavus*, *Aspergillus niger*, *Epidermophyton floccosum*, *Trycophyton*

mentagrophytes and *Microsporium canis*. Two strains of yeast have been investigated in this study; *Candida albicans* and *Cryptococcus neoformans*

Preparation of plant extracts

Organic solvents used in this experiment were obtained from Sigma-Aldrich Company. The dried roots, seeds or leaves were ground into fine powder with an electric blender. Fifty grams were suspended in hot water or organic solvents (methanol, diethyl-ether, ethyl acetate and chloroform) in sterile 250ml conical flasks and kept at 4°C overnight. After overnight incubation, the supernatant was filtered through **Whatman** No.1 filter paper and the filtrate was concentrated by evaporation in a rotary evaporator at 40°C. The residue was weighed, dissolved in 5% dimethyl sulfoxide (DMSO) and stored in the refrigerator at 4°C prior to use.

Antimicrobial Activity

This test was carried out using agar well diffusion method according to Joshi *et al.* (2009). Bacteria or yeast were taken and shaken in the sterile distilled water corresponding to 10^8 CFU/ml for bacteria and 10^6 CFU/ml of yeast (Mihajilov-Krstev *et al.*, 2010). Fungal inocula were prepared by flooding Petri dish with 8 to 10 ml of distilled water and the conidia were scraped using sterile spatula. The spore density of each fungus was 1.5×10^4 spore/ml (Adigüzel *et al.*, 2005). Minimal inhibitory concentration was determined by the method recommended by Ter-Laak *et al.* (1991). Each antimicrobial agent was serially diluted by transferring 100 µl of the antimicrobial agent through sterilized microtitre plate containing 100 µl media (nutrient broth for bacteria, Sabouraud dextrose broth for fungi and yeast). Freshly

prepared standard number of cells (1.5×10^8 CFU / ml for bacteria or yeast and 4×10^4 spore/ml for fungal isolates) was added to the media that contained some drops of phenol red. Glucose metabolisms were measured by a change of the color of phenol red indicator from red to yellow. MIC was determined at the concentration with no color change and DMSO was used as a control.

Toxicity and antitumor activity of the plant extracts

Cytotoxicity is performed by Brine Shrimp Test (BST). Different dilutions of each plant extract made in DMSO were prepared (100, 200, 300 and 400 $\mu\text{g/ml}$) and 0.5 ml of each dilution was added to 4.5 ml of brine solution and maintained at room temperature for 24 h under the light (Krishnaraju *et al.*, 2005). The surviving larvae were counted by light microscope and the number of dead shrimps in each vial was recorded. The concentration at which 50% of the larvae were killed (LD_{50}) was determined as the toxic concentration (Lachumy *et al.*, 2010). The antitumor activity of the tested plants was determined against Ehrlich carcinoma and Lymphoma cell line. Cells were grown in RPMI 1640 medium (Sigma, USA) with 10% fetal calf serum (FCS) (Gibco, USA) at 37°C under a humidified atmosphere consisting of 95% air and 5% CO_2 for 48 h. The percentage of cell viability was assessed to determine the 50 % lethal dose by which 50% of cells are killed (LD_{50}).

RESULTS AND DISCUSSION

The non-availability and high cost of new generation antibiotics with limited effective span have resulted in increase in morbidity and mortality (Williams, 2000). Therefore, there is a need to look for substances from other sources with proven antimicrobial activity. Consequently, this has led to the search for more effective antimicrobial agents among materials of plant origin, with the aim of discovering potentially useful

active ingredients that can serve as source and template for the synthesis of new antimicrobial drugs (Pretorius *et al.*, 2003).

In this study, three plants were collected and extracted using hot water, methanol, ethyl acetate, N-butanol, diethyl ether and chloroform. The tested plants were *S. incanum*, *E. cardamomum* and *Z. officinale*.

The tested plants were selected based on traditional medicine knowledge used by Saudi Arabian people. All the obtained extracts were screened for their antibacterial activity against *E. coli*.

It was clear that all of the extracts have inhibitory activity against *E. coli*. The inhibitory activity varied according to the solvent used. The methanol extracts for all of tested plants showed significant antibacterial activities against *E. coli* compared to the activity of the other solvents which almost have similar effect. Previous studies provide similar results for methanol as a better solvent for more consistent extraction of antimicrobial substances from medicinal plants as compared to other solvents such as water and ethanol (Ahmad *et al.*, 1998), hot water, N- butanol, diethyl-ether, ethyl acetate and chloroform (El Sayed and Aly, 2014).

Since methanol extract was the most active solvent, its activity was tested against the other pathogenic organisms including Gram positive bacteria (*S. aureus*, *S. aureus* MRSA and *E. faecalis* VRE), Gram negative bacteria (*P. aeruginosa*, *K. pneumonia*, *A. baumannii* and *P. mirabilis*), yeast (*C. albicans*, and *C. neoformans*) and molds (*A. flavus*, *A. niger*, *E. floccosum*, *T. mentagrophytes* and *M. canis*) (Table 3, 4).

S. incanum has an effect on Gram negative and Gram positive isolates, but zones of inhibition are greater in Gram negative organisms (17-29 mm.). It has the best activity

against *P. aeruginosa*. MIC values were 50-150 µg/ml. The most affected Gram negative organism was *E. coli*, while *E. faecalis* was the most resistant organism.

In accordance, Omwenga *et al.* (2012) found that the methanolic extract of *S. incanum* has better activity on *P. aeruginosa* producing a wide zone of inhibition compared to *S. aureus* and *B. subtilis*. Such results are promising since *P. aeruginosa* is hard to be controlled by most antibiotics due to its cell wall properties (Omwenga *et al.*, 2009).

Z. officinale is exceeding the effect of *S. incanum* against Gram negative bacteria (inhibition zones, 22-24 mm., MIC 50-100 µg/ml.). Significant antimicrobial activity of *Z. officinale* against *E. coli* has been reported by Yahaya *et al.* (2012). Gull *et al.* (2012) reported antimicrobial activity of the plant methanolic extract on both Gram positive and Gram negative bacteria. The antibacterial activities of the extracts are expected perhaps due to the compounds like flavonoids and volatile oil which were dissolved in organic solvents. It is reported that sesquiterpenoids are the main component of ginger which attributes its antibacterial activity (Malu *et al.*, 2008)

The greater antimicrobial activity for the methanolic extracts of the three tested plants was *E. cardamomum*. It has relative increasing in inhibition zones compared to *Z. officinale* with exception of *K. pneumonia*, *P. mirabilis* and *S. aureus* 33591 as *Zingiber officinale* had a stronger effect.

The inhibitory effect on Gram negative strains was greater than Gram positives. The markedly increase in the antimicrobial activity against *P. aeruginosa*, Gram negative bacteria and *E. faecalis* (VRE) among Gram positive bacteria is contributing to the production of drugs effective against multi-drug resistant bacteria. In accordance, Jemal *et al.* (2011) observed that complete (100%) growth inhibition was at 15% cardamom hydrosol concentration against *E. coli*, *S. aureus* and *S. typhi*, but in contrast, he found *P. aeruginosa* as the most resistant organism.

Antifungal activity of the selected plants has also been investigated and represented in Tables (5) and (6). *E. cardamomum* had the highest values of inhibition on filamentous fungi with exception of *A. niger* compared to methanolic extracts of other plants. *S. incanum* has the highest effect on such strain. Inhibition zones of *E. cardamomum* ranged from 11 mm. to 17 mm. the former was recorded for *A. niger*. MIC were 50-100 µg/ml. Obvious antimicrobial action on dermatophytes (*E. floccosum*, *T. mentagrophytes* and *M. canis*) was observed. It also recorded considerable effect against *C. albicans*. A similar result was obtained by Ağaoğlu *et al.* (2005). According to Aneja and Sharma (2010), *E. cardamomum* displayed good to moderate activity against *C. albicans*. *E. cardamomum* is widely used in various parts of the world's traditional medicine system and it has been used in India since ancient times (Dhulap *et al.*, 2008). The antimicrobial potential of this plant extracted in different solvents (e.g. aqueous, methanol, ethanol, acetone, chloroform, hexane, ethyl acetate and diethyl ether) had been evaluated against different bacterial and fungal human pathogens and had reported variable activities in different parts, seeds, pods and fruits in different solvents (Dhulap *et al.*, 2008; Agaoglu *et al.*, 2005; Kaushik *et al.*, 2010; Singh *et al.*, 2008; Aneja and Joshi, 2009).

Cytotoxic effect using brine shrimp and antitumor activity using Erlich cell line and lymphoma cell line of the plants under investigation were studied and represented in Table (7). The highest cytotoxic effect was recorded for the methanolic extract of *S. incanum* (LD₅₀ is 600), while the minimal effect was recorded for *E. cardamomum* (LD₅₀ is ≥ 600). Antitumor activity of *E. cardamomum*, and *Z. officinale* showed LD₅₀ ≥ 600 µg/ml on both Lymphoma cell line and Erlich cell line, while it was 400 µg/ml on *S. incanum* µg/ml on both Erlich cell line and lymphoma cell line.

CONCLUSION

On the basis of the experimental results and discussion, it can be postulated that methanol is the best solvent for all of the plants under investigation. *S. incanum* and *Z. officinale* methanolic extracts can be used effectively against *E. coli*. Moreover, *S. incanum* was the best plant extract against *P. aeruginosa*. Methanolic extract of *Z. officinale* provides the best action against *S. aureus* (MRSA) and *K. pneumonia* (ESBL). So, it is considered to be as promising antibiotic against multidrug resistant bacteria. *E. cardamomum* has broad spectrum antimicrobial activity against Gram negative bacteria, Gram positive bacteria, yeast and filamentous fungi.

REFERENCES

- Adigüzel, A.; Güllüce, M.; Şenqüel, M.; Öğütcü, H.; Şahian, F. and Karaman, I. (2005) Antimicrobial effects of *Ocimum basilicum* (Labiatae) extract, Turkish J. Biotechnol., 29: 155-160.
- Agaoglu S; Dostbil N; Alemdar S. (2005). Antimicrobial effect of seed extract of Cardamom (*Elettaria cardamomum* Maton). YYU Vet Fak Derg, 16 (2):99-101.
- Ahmad I; Mehmood Z. and Mohammad F. (1998). Screening of some Indian medicinal plants for their antimicrobial properties. Journal of Ethnopharmacology; 62:183-193. Available: [http://dx.doi.org/10.1016/S0378-8741\(98\)00055-5](http://dx.doi.org/10.1016/S0378-8741(98)00055-5).
- Aibinu, I; Adenipekun, E; and Odugbemi, T. (2004). Emergence of quinolone resistance amongst *Escherichia coli* strains isolated from clinical infections in some Lagos State Hospitals in Nigeria. Nigerian Journal of Health and Biomedical Science, 3(2):73-78.
- Aibinu, I. E.; Ohaegbulam, V. C.; Adenipekun, E. A.; Ogunsola, F. T.; Odugbemi, T. O. and Mee, B. J. (2003). Extended-spectrum beta-lactamase enzymes in clinical isolates of *Enterobacter* species from Lagos, Nigeria. Journal of Clinical Microbiology. **41(5)**:2197-2200.
- Aneja K.R. and Radhika Joshi (2009). Antimicrobial activity of *Amomum subulatum* and *Elettaria cardamomum* against dental caries causing microorganisms. Ethnobotanical Leaflets 13: 840-49. Corresponding Author: joshi_radhika31282@yahoo.com
- Aneja K.R. and Chetan Sharma (2010). Antimicrobial potential of fruit extracts of *Elettaria cardamomum* Maton (CHHOTI ELAICHI) against the pathogens causing ear infection. Pharmacologyonline 3: 750-756.
- Bari, M.A.; Islam, W.; Khan A.R. and Abul Mandal. (2010). Antibacterial and antifungal activity of *Solanum torvum* (Solanaceae). International Journal of Agricultural journal .12 (3): 386-390.

Bartley J. and Jacobs A. (2000). Effect of drying on flavor compounds in Australian-grown ginger (*Zingiber officinale*). Journal of the Science of Food and Agriculture. 80:209-215.

Britto SJ. and Senthinkumar S. (2001). Antimicrobial activities of *Solanum incanum* leaf extract. Asian J, Microbio Biotech and EnviroSci, 3(1-2):65-66.

Davis J. (1995). Inactivation of the antibiotics and the dissemination of resistance genes. Science 264: 375-382, 1994). (Service RF, Antibiotics that resist resistance. Science 270: 724-727, 1995)

Denston, T.C. (1951). A Textbook of Pharmacology, 5th edition, Medicinal Plants of West Africa. London Oxford Press. 147 – 149.

Dhulap S; Anita M. and Hirwani RR. (2008). Phyto-pharmacology of *Elettaria cardamomum*. Phcog Rev Sup 2(4): 27-35.

El Sayed H. and Aly M. (2014). Antibacterial activities of six medicinal plants used traditionally by Saudi people to treat common diseases. British Biotechnology Journal, 4(4): 499-510.

Gull I.; Mariam Saeed; Halima Shaukat; Shahbaz M Aslam; Zahoor Qadir Samra and Amin M Athar (2012). Inhibitory effect of *Allium sativum* and *Zingiber officinale* extracts on clinically important drug resistant pathogenic bacteria. Annals of Clinical Microbiology and Antimicrobials 11:8-22.

Imadia K.; Fukushima S.; Shirai T.; Ohtani M.; Nakanish K. and Ito N. (1983). Promoting activities of butylated hydroxytoluene on 2-stage urinary bladed carcinogenesis and inhibition of γ -glutamyl transpeptidase development in the liver of rats. Carcinogenesis. 4:895-899.

Jemal Hussien; Chalachew Teshale and Jemal Mohammed (2011). Assessment of the antimicrobial effects of some ethiopian aromatic spice and herb hydrosols. International Journal of Pharmacology, 7 (5): 635-640.

Joshi, B.; Lekhak, S. and Sharma, A. (2009): Antibacterial property of different medicinal plants: *Ocimum sanctum*, *Cinnamomum zeylanicum*, *Xanthoxylum armatum* and *Origanum majorana*. Journal of Science, Engineering and Technolgy, 5 (I): 143- 150.

Kapoor L. D. (2000). Handbook of Ayurvedic Medicinal Plants. CRC Press, Boca Raton, FL.

Kaushik P; Goyal P; Chauhan A and Chauhan G. (2010) *In Vitro* evaluation of antibacterial potential of dry fruit extracts of *Elettaria cardamomum* Maton (Chhoti Elaichi). Iranian J Pharm Res, 9 (3): 287-292.

Krishnaraju, A.V.; Rao, T.V.N.; Sundararaju, D.; Vanisree, M.; Tsay, H. and Subbaraju, G. V. (2005): Assessment of bioactivity of Indian medicinal plants using brine shrimp (*Artemia salina*) lethality assay. International Journal of Applied Science and Engineering. 3, 2: 125-134.

Lachumy, S. J.; Zuraini Z. and Sasidharan, S. (2010): Antimicrobial activity and toxicity of methanol extract of *Cassia fistula* seeds. Research Journal of Pharmaceutical, Biological 398. and Chemical Sciences, 1(4): 391-

Malu SP; Obochi GO; Tawo EN and Nyong BE (2008). Antibacterial activity and medicinal properties of ginger (*Zingiber officinale*). Global J Pure Appl Sci 2008, 15:365–368.

Martini, N. and Eloff, J.N. 1998. The preliminary isolation of several antibacterial compounds from *Combretum erythrophyllum* (Combretaceae). Journal of Ethnopharmacology, 62:255-2643.

McGee H. (2004). On Food and Cooking: The Science And Lore Of The Kitchen (2nd ed.). New York: Scribner.425-426.

McKenna, J. 1996. Alternatives to Antibiotics. 55-58.

Mihajilov-Krstev, T.; Radnović, D. and Kitić, D. (2010) Antimicrobial activity of Satureja L. essential oils against phytopathogenic bacteria *Erwinia amylovora*, *Biologica Nyssana*., 1 (1-2): 95-98

Miri P; Jungdon B and Dae-Sil L. (2008). Antibacterial activity of [10]-gingerol and [12]-gingerol isolated from ginger rhizome against periodontal bacteria. *Phytotherapy Research*, 22(11):1446–1449.

Omwenga E.O.; Ogol Calistus; Paul Kamau Mbugua and Paul Owour Okemo. (2012). Ethnobotanical identification and anti-microbial evaluation of some anti-diarrhoeal plants used by the Samburu Community, Kenya. *Malaysian Journal of Microbiology*, 8(2). 68-74.

Omwenga E.O; Okemo P.O; Mbugua P.K, and Ogol C.K.P, (2009). Ethnobotanical survey and the antimicrobial evaluation of medicinal plants used by the Samburu community (Kenya) for treatment of diarrhoea. *Pharmacognosy Magazine*, 4 (18): 165-176.

Oyagbemi AA; Saba Aba nd Azeez OI. (2010). Molecular targets of [6]-gingerol: Its potential roles in cancer chemoprevention. *Biofactors*, 36(3):169-178.

Park M; Bae J and Lee D. (2008). Antibacterial activity of [10]-gingerol and [12]-gingerol isolated from ginger rhizome against periodontal bacteria. *Phytother. Res.*, 22:1446-1449.

Patil S.; Joshi Vijaya, Sutar Prasanna and Sambrekar Sudhir, (2009). Screening of whole plant extract of *Solanum surattense* for antibacterial activity. *International Journal of Pharmaceutical Sciences*, 1(1):110-114.

Pavitra P.S.; Janani V.S.; Charumathi K.H.; Indumathy R.; Sirisha Potala and Rama S. Verma (2012). Antibacterial activity of plants used in Indian herbal medicine. *International Journal of Green Pharmacy*, 23-28.

Pretorius, J.C.; Magama S., and Zietsman P.C. (2003). Growth inhibition of plant pathogenic bacteria and fungi by extracts from selected South African plant species *South African Journal of Botany*, 20: 188-192.

Schelz Z; Molnar J. and Hohmann J. (2006). Antimicrobial and antiplasmid activities of essential oils. *Fitoterapia.*,77(4):279-85.

- Sheeba, E. (2010). Antibacterial activity of *Solanum surattense* Burm. F. Kathamandu University Journal of Science, Engineering and Technology. 6: (I). 1-4.
- Sieradzki K.; Roberts R.B.; Haber S.W. and Tomasz A. (1999): The development of vancomycin resistance in a patient with methicillin-resistant *Staphylococcus aureus* infection. N. Engl. J. Med., **340**: 517–523.
- Singh G; Kiran S; Marimuthu P; Isidorov V. and Vinogorova V. (2008). Antioxidant and antimicrobial activities of essential oil and various oleoresins of *Elettaria cardamomum* (seeds and pods). J Sci Food Agric., 88:280–289.
- Soriful I.; Atikur R.; Mominul I. S.; Mostafizur RAHMAN; Abu Hena Mostofa JAMAL and Firoz ALAM (2010). *In vitro* antibacterial activity of methanol seed extract of *Elettaria cardamomum* (L.) Maton. Agriculturae Conspectus Scientificus., 75 (3). 113-117.
- Sridhar T.M.; Josthna P.; and Naidu C.V.; (2011). *In vitro* Antibacterial activity and phytochemical analysis of *Solanum nigrum* (Linn.) – An important antiulcer medicinal plant. Journal of Experimental Sciences, 2(8): 24-29.
- Ter-Laak E.A.; Pijpers A.; Noordergraaf; J.H.; Schoevers E.C. and Verheijden, J.H. (1991) Comparison of methods for *in vitro* testing of susceptibility of porcine *Mycoplasma* species to antimicrobial agents. Antimicrob Agents Chemother **35**, 228–233.
- Vargas AC; Loguercio AP; Witt NM; Costa MM; Silva MS and Viana LR. (2004). Alcoholic propolis extract: antimicrobial activity. Cienc Rural., 34(1):159-63.
- Williams, R., (2000) Antimicrobial resistance a global threat. Essential Drug Monitor, 1: 28-29
- Wood GB. (2013). A treatise on therapeutics, and pharmacology or materia medica: Volume 2. J. B. Lippincott & Co. 1867a. Retrieved 2 March; 2013.
- World Health Organization (1978): The promotion and development of traditional medicine. Technical report series, 622 pp.

Yahaya O.; Yabefa, J.A.; Umar I.O.; Datshen M.M; Egbunu Z.K. and Ameh J. (2012). Combine antimicrobial effect of ginger and honey on some human pathogens. British Journal of Pharmacology and Toxicology 3(5): 237-239.

Table (1). The selected medicinal plants, their families and common names

Scientific name	Family	Common name	Used part
<i>Solanum incanum</i>	Solanaceae	Bitter apple	fruits
<i>Zingiber officinale</i>	Zingiberaceae	Ginger	Root
<i>Elettaria cardamomum</i>	<u>Zingiberaceae</u>	Cardamom	fruits

Table (2): The antibacterial activity of some plants extracts using water and organic solvents against *E. coli*.

Diameter of inhibition zone mm						
Solvent Plant	Hot water	Methanol	N- butanol	Diethyl- ether	Ethyl acetate	Chloroform
<i>S. incanum</i>	19 ± 0.07	39 ± 1.0	19 ± 0.9	17 ± 0.2	15 ± 0.9	18 ± 0.4
<i>Z. officinale</i>	18 ± 0.17	39 ± 1.0	19 ± 0.6	17 ± 0.2	15 ± 1.0	18 ± 2.0
<i>E. cardamomum</i>	17±1.30	26 ± 1.0	18 ± 0.3	17 ± 0.4	19 ± 2.0	13 ± 1.0

Bacterial index*	54	104	56	51	49	49
------------------	----	-----	----	----	----	----

*Bacterial Index: Total Activity (mm).

Table (3). Antibacterial Activities (Diameter of the Inhibition Zone, mm) of the methanolic plant extracts against different pathogenic bacteria

Plants Bacteria	<i>S. incanum</i>	<i>E. cardamomum</i>	<i>Z. officinale</i>	Control (DMSO)
<i>P. aeruginosa</i>	29±1.0	27±0.2	23±3.0	ND
<i>K. pneumoniae</i>	16±1.0	16±0.2	22±2.8	ND
<i>P. mirabilis</i>	17±2.1	16±0.4	24±0.5	ND
<i>A. baumannii</i>	19±0.8	26±0.4	12±0.6	ND
<i>E. faecalis</i>	11±0.5	14±0.6	11±0.4	ND
<i>S. aureus</i> 33591	14±0.9	12±0.4	18±0.4	ND
<i>S. aureus</i> 25923	14±1.0	18±0.8	14±0.8	ND
Bacterial index	120	129	124	ND

*Bacterial Index: Total Activity (mm).

Table (4). The minimal inhibitory concentration (µg/ml) of methanolic extracts against different pathogenic bacteria

Plants Bacteria	<i>S. incanum</i>	<i>E. cardamomum</i>	<i>Z. officinale</i>	Control (DMSO)
<i>P. aeruginosa</i>	50	50	100	> 200
<i>K. pneumoniae</i>	150	150	50	> 200
<i>P. mirabilis</i>	150	100	50	> 200
<i>A. baumannii</i>	150	50	150	> 200
<i>E. faecalis</i>	150	150	150	> 200
<i>S. aureus</i> 33591	100	150	100	> 200

<i>S. aureus</i> 25923	100	100	100	> 200
<i>E.coli</i>	100	50	100	> 200

Table (5): Effect of the methanolic extracts on fungi and yeast

*Fungal Index: Total Activity (mm).

Table (6). The minimal inhibitory concentration ($\mu\text{g/ml}$) of methanolic extracts against different pathogenic fungi and yeast, compared with that of control

Plants Fungi	<i>S. incanum</i>	<i>E. .cardamomum</i>	<i>Z. officinale</i>	Control (DMSO)
<i>A. flavus</i>	50	100	100	>200
<i>A. niger</i>	50	100	100	>200
<i>E. floccosum</i>	100	100	50	>200
<i>T. mentagrophytes</i>	100	50	50	>200

Plants Fungi	<i>S. incanum</i>	<i>E. .cardamomum</i>	<i>Z. officinale</i>	Control (DMSO)
<i>A. flavus</i>	11 \pm 1.0	17 \pm 0.23	13 \pm 1.3	ND
<i>A. niger</i>	17 \pm 0.8	11 \pm 0.08	10 \pm 0.5	ND
<i>E. floccosum</i>	11 \pm 1.2	14 \pm 0.22	11 \pm 0.8	ND
<i>T.mentagrophytes</i>	11 \pm 1.1	13 \pm 0.43	11 \pm 1.5	ND
<i>M. canis</i>	11 \pm 0.18	16 \pm 0.14	11 \pm 0.16	ND
<i>C.albicans</i>	10 \pm 0.15	16 \pm 0.16	11 \pm 0.10	ND
<i>C. neoformans</i>	13 \pm 1.4	11 \pm 0.8	13 \pm 0.18	ND
Fungal index*	84	98	80	ND

<i>M. canis</i>	50	100	50	>200
<i>C. albicans</i>	50	50	100	>200
<i>C. neoformans</i>	150	150	150	>200

Table (7). Toxicity against *Artimia salina* (% mortality) and antitumor activities of the different concentrations of plant methanolic extracts

Plants	Toxicity against <i>Artimia salina</i> (% of mortality) at different concentrations ($\mu\text{g/ml}$)					Antitumor activity (LD_{50} , $\mu\text{g/ml}$)	
	Control	200	400	600	LD_{50}	Lymphoma cell line	Erlich cell line
<i>S. incanum</i>	0	10	20	59	500	400	400
<i>E. cardamomum</i>	0	0	0	10	≥ 600	≥ 600	≥ 600
<i>Z. officinale</i>	0	0	10	26	≥ 600	≥ 600	≥ 600